

家畜衛生

(1) 家畜伝染病発生状況 ※患畜のみ

| 年次 | | 51~57年 | 60 | 2 | 7 | 12 | 17 | 21 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | R1 | R2 | R3 | R4 | R5 | R6 | R7 |
|-----|-----------------|--------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| 畜種 | 病名 | | | | | | | | | | | | | | | | | | | | | | |
| 牛 | 流行性脳炎(頭) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | ブルセラ病(頭) | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 結核病(頭) | 35 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 炭疽(頭) | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | ヨーネ病(頭) | 0 | 0 | 0 | 0 | 10 | 20 | 5 | 5 | 2 | 8 | 8 | 8 | 4 | 1 | 0 | 2 | 0 | 0 | 1 | 4 | 3 | 3 |
| 馬 | 馬伝染性貧血(頭) | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 豚 | 流行性脳炎(頭) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 1 | 0 | 0 | 0 |
| | 豚熱(頭) | 369 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 45 | 0 | 0 | 0 | 0 | 0 | 0 |
| 鶏 | 高病原性鳥インフルエンザ(羽) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0 | 0 | 19 |
| うずら | 高病原性鳥インフルエンザ(羽) | 0 | 0 | 0 | 0 | 0 | 0 | 40 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 蜜蜂 | 腐蛆病(群) | 502 | 64 | 39 | 12 | 0 | 1 | 0 | 1 | 5 | 15 | 22 | 9 | 12 | 3 | 13 | 13 | 5 | 14 | 0 | 0 | 1 | 0 |